



Nicorette QuickMist SmartTrack Mouthspray

Uses

Relieves and/or prevents craving and nicotine withdrawal symptoms associated with tobacco dependence. It is indicated to aid smokers wishing to quit or reduce prior to quitting, to assist smokers who are unwilling or unable to smoke, and as a safer alternative to smoking for smokers and those around them. It is indicated in pregnant and lactating women making a quit attempt.

Dosage

Over 12 years: the patient should make every effort to stop smoking or vaping completely during treatment with Nicorette QuickMist. One or two sprays used when normally would smoke or vape or if cravings emerge. Maximum dose of 4 sprays per hour, not exceeding 2 sprays per dosing episode, and 64 sprays in any 24-hour period. Nicorette QuickMist should be used whenever the urge to smoke or vape is felt or to prevent cravings in situations where these are likely to occur. Patients able to stop smoking/vaping immediately should initially replace all their cigarettes/e-cigarettes with the Nicorette QuickMist and as soon as they are able, reduce the number of sprays used until they have stopped completely. Patients aiming to reduce cigarettes/e-cigarettes should use the mouthspray between smoking/vaping episodes, as needed, to prolong smoke/vape-free intervals and to reduce their use as much as possible. As soon as they are ready patients should aim to quit smoking/vaping completely. Under 12 years: not to be used.

Format and Ingredients

Mouthspray containing:
Nicotine 1mg,

