Corsodyl 1%w/w Dental Gel

Uses

For the inhibition of formation of dental plaque and as an aid in the treatment and prevention of gingivitis and maintenance of oral hygiene. For use in a post-peridontal surgery or treatment regimen to promote gingival healing. Useful in the management of recurrent mouth ulcers and recurrent oral candidal infections (e.g denture stomatitis and thrush) and as an aid in the prevention of dental caries in high caries-risk patients (e.g. xerostomia sufferers) in a regimen as an adjunct to fluoride.

Dosage

Adults, the elderly and children over 12 years: brush teeth with 1 inch of the gel on a moistened toothbrush, once or twice daily for about 1 minute. For gingivitis, use for 1 month. For mouth ulcers and oral candidal infections: apply once or twice daily for about 1 minute. When used as part of a regimen to prevent dental caries, apply for 5 minutes nightly for 14 days in a close fitting tray. Children under 12 years: should not use unless recommended by a healthcare professional.

Format and Ingredients

Gel containing:
Chlorhexidine Digluconate 1% w/w,

Category:
Mouth Care

Manufacturer:
HALEON PLC

Pack size: 50g

RRP: £7.25

Legal Status: (P)

A division of GROUPE EUROCOM
PAGB
Communications International Group

Product data © Communications International Group

Communications International Group Ltd 2026,
Linen Hall, 162-168 Regent Street, London W1B 5TB