



NiQuitin Mint 2mg, 4mg Lozenges

Uses

For the treatment of tobacco dependence by relieving nicotine withdrawal symptoms including cravings associated with smoking cessation. Should be used with behavioural support. NiQuitin Mint 2mg Lozenge: for those who smoke more than 30 minutes after waking. NiQuitin Mint 4mg Lozenge: for those who smoke within 30 minutes of waking.

Dosage

Lozenges should be used whenever there is an urge to smoke. Adult smokers 18 years and over: weeks 1 to 6: at least 9 lozenges but no more than 15 lozenges per day. Gradually reduce use so that by week 12, 1 to 2 lozenges a day are used only on occasions when strongly tempted to smoke. A healthcare professional should be consulted if unable to make a permanent quit attempt after 6 months. Allow lozenge to dissolve in the mouth, occasionally moving lozenge from one side to the other, until dissolved (about 20-30 minutes). Do not chew or swallow whole. Adolescents (12-17 years): should use the lozenges only on the advice of a healthcare professional. Under 12 years: not recommended.

Format and Ingredients

Cream/white, round, biconvex lozenge containing: Nicotine 2 or 4mg,

Category:
Smoking Cessation

Manufacturer:
PAGB

Pack size: Both strengths:
72 lozenges, 132 lozenges

RRP: £10.00, £14.99