



Nicorette Lozenge

Uses

Relieves and/or prevents cravings and nicotine withdrawal symptoms associated with tobacco dependence. It is indicated to aid smokers wishing to quit or reduce prior to quitting, to assist smokers who are unwilling or unable to smoke, and as a safer alternative to smoking for smokers and those around them. It is indicated in pregnant and lactating women making a quit attempt.

Dosage

12 years and over: The strength of lozenge to be used will depend on the smoking habits of the individual. If the patient smokes 20 or fewer cigarettes a day, 2mg nicotine lozenge is indicated. If more than 20 cigarettes per day are smoked, 4mg nicotine lozenge should be used. The lozenges should be used whenever the urge to smoke is felt or to prevent cravings in situations where these are likely to occur. Smokers willing or able to stop smoking immediately should initially replace all their cigarettes with the lozenge and as soon as they are able, reduce the number of lozenges used until they have stopped completely. Smokers aiming to reduce cigarettes should take the lozenge, as needed, between smoking episodes to prolong smoke-free intervals and with the intention to reduce smoking as much as possible. As soon as they are ready smokers should aim to quit smoking completely. Most smokers require 8 to 12 lozenges per day, not to exceed 15 lozenges. When making a quit attempt behavioural therapy, advice and support will normally improve the success rate. Those who have quit smoking but are having difficulty discontinuing with the lozenge are recommended to contact their pharmacist or doctor for advice.

Format and Ingredients

Lozenge containing:
Nicotine 2mg, Nicotine 4mg,

Manufacturer:
KENVUE

Pack size: Mint, Fruit 2mg:
40, Mint, Fruit 2mg: 80, Mint,
Fruit 2mg: 160, Mint 4mg: 40,
Mint 4mg: 80

RRP: £9.52, £17.14, £32.38,
£9.52, £17.14

Legal Status: (GSL), (GSL),
(GSL), (GSL), (GSL)

A division of
 GROUPE
EUROCOM

 PAGB

 **Communications
International
Group**