NiQuitin Mint 2mg, 4mg Lozenges

Uses

For relief of nicotine withdrawal symptoms including cravings associated with smoking cessation. 2mg Lozenge: for those who smoke more than 30 minutes after waking. 4mg Lozenge: for those who smoke within 30 minutes of waking.

Dosage

Lozenges should be used whenever there is an urge to smoke. Adult smokers 18 years and over: weeks 1-6: at least 9 lozenges but no more than 15 lozenges a day. Gradually reduce so that by week 12, 1-2 lozenges a day are used only on occasions when strongly tempted to smoke. A healthcare professional should be consulted if unable to make a permanent quit attempt after 6 months. Allow lozenge to dissolve in the mouth, occasionally moving it from one side to the other until dissolved (about 20-30 minutes). Do not chew or swallow whole. Adolescents (12-17 years): use only on the advice of a healthcare professional. Under 12 years: not recommended.

Format and Ingredients

Lozenge containing:
Nicotine 2 or 4mg,

Category:
Smoking Cessation

Manufacturer:
PERRIGO

Pack size: Both strengths: 72 lozenges, 132 lozenges

RRP: £10.00, £14.99

Legal Status: (GSL),

A division of GROUPE EUROCOM
PAGB
Communications International Group

Communications International Group Ltd 2026,
Linen Hall, 162-168 Regent Street, London W1B 5TB

Product data © Communications International Group