Canesten Thrush Duo Oral Capsule & External Cream

Uses

Canesten Thrush Oral Capsule can be used to treat acute or recurrent candidal vaginitis. It should also be used for the treatment of partners with associated candidal balanitis. Canesten Thrush External Cream is indicated for the treatment of candida vulvitis. It should be used as an adjunct to treatment of candidal vaginitis. It can also be used for treatment of the sexual partner's penis to prevent re-infection.

Dosage

16-60 years: 1 capsule should be swallowed whole. The cream should be applied thinly 2 or 3 times daily to the vulva and surrounding area and rubbed in gently. Treatment with the cream should be continued until symptoms of the infection disappear. However, if after concomitant treatment of the vaginitis, symptoms do not improve within 7 days, the patient should consult a doctor. If the cream is being used for treatment of the sexual partner's penis, it should be applied 2 or 3 times daily for up to 2 weeks. Under 16 or over 60: a doctor should be consulted.

Format and Ingredients

Oral capsule and tube of external cream containing: Capsule: Fluconazole 150mg, External Cream: Clotrimazole 2% w/w,

Category:
Women's health

Manufacturer:

Pack size: Pack

RRP: £19.69

A division of GROUPE EUROCOM
PAGB
Communications International Group

Product data © Communications International Group

Communications International Group Ltd 2026,
Linen Hall, 162-168 Regent Street, London W1B 5TB