Piriton Syrup

Uses

For symptomatic control of all allergic conditions responsive to antihistamines, including hayfever, urticaria, food allergy and insect bites. Can also be used to relieve the itch associated with chickenpox.

Dosage

Adults and children 12 years and over: 10ml (4mg) every 4-6 hours. Maximum daily dose: 60ml (24mg). Elderly: consider using a lower daily dose (e.g. maximum 12mg). 6-12 years: 5ml (2mg) every 4-6 hours. Maximum daily dose: 30ml (12mg). 2-6 years: 2.5ml (1mg) every 4 to 6 hours. Maximum 15ml (6mg) in any 24 hours. Infants 1-2 years: 2.5ml (1mg) twice daily. Maximum daily dose: 5ml (2mg). Under 1 year: not recommended. The minimum interval between the doses should be 4 hours.

Format and Ingredients

Syrup containing per 5ml:
Chlorphenamine Maleate 2mg,

Category:
Allergies, Babies & children

Manufacturer:
HALEON PLC

Pack size: 150ml

RRP: £6.50

Legal Status: (P)

A division of GROUPE EUROCOM
PAGB
Communications International Group

Product data © Communications International Group

Communications International Group Ltd 2026,
Linen Hall, 162-168 Regent Street, London W1B 5TB